

Thyroid Lesions: A Unique Approach

Som Biswas^{1*} and Srirupa Biswas²

¹Resident, Department of Radio diagnosis, GRMC, Gwalior

²Assistant Professor, School of Pharmacy, ITM University, Gwalior

***Corresponding Author:** Dr Som Biswas, Resident, Department of Radio diagnosis, GRMC, Gwalior, India.

Received: November 18, 2017; **Published:** November 29, 2017

Volume 1 Issue 1 November 2017

© All Copy Rights are Reserved by Som Biswas and Srirupa Biswas.

Introduction

Thyroid lesions are a common finding in the general population, especially in iodine deficiency areas. With the more radical use of iodized salt and other measures the incidence of earlier common disorders like goiter, congenital hypothyroidism, etc. have dropped dramatically but still these are quite common in developing countries. Among the several imaging techniques that provide clinically useful anatomic information and pathologic condition about thyroid gland, sonography has become the method that is most commonly employed. Its use has resulted in early and accurate detection of various thyroid disorders. This is of immense help in timely management and prevention of complications. Colour Doppler Sonography is used in thyroid vascular study. Dynamic information such as velocity and direction of blood flow as well as degree of vascularity of organ can be revealed by colour doppler studies. Ultrasound vascular study is a noninvasive and low cost method

A study of thyroid lesion was done in 100 patients in leading teaching hospital, an attempt was made to evaluate and establish certain gray scale and colour Doppler criteria for distinguishing between benign and malignant thyroid lesions.

Aims & Objectives

1. Evaluation of common thyroid disorder in our setup.
2. To evaluate the accuracy of ultrasound in diagnosing thyroid disorders.
3. To evaluate the important sonological parameters in various thyroid disease.
4. To differentiate between solid, cystic and mixed nodular?
5. To differentiate between benign and malignant thyroid lesions.
6. To evaluate vascular nature of various thyroid lesion using colour Doppler.
7. Role of colour Doppler in distinguishing benign for malignant thyroid nodules.
8. Comparison of clinical, ultrasound colour Doppler and FNAC (fine needle aspiration cytology) findings.

Material and Methods

Patient selection & clinical presentation

Random selection of patient of all age groups was done. They presented with sign and symptoms suggestive of thyroid disorders.

Citation: Som Biswas and Srirupa Biswas. "Thyroid Lesions: A Unique Approach". *Chronicle of Medicine and Surgery* 1.1 (2017): 10-17.

The commonest clinical symptom was swelling in front of neck, which moves up with deglutition. There were systemic symptoms like pain and fever in thyroiditis weight loss and palpitation in hyperthyroidism, weight gain and hoarseness of voice in hypothyroidism and rapid increase in size with weight loss in malignancies. In nodular goiter, patients were usually thyroid. The nodules were palpable and sometimes visible.

The study was done from October 2013 to September 2014 in the Tertiary Hospital. 100 patients underwent sonography of the thyroid.

Technical consideration

- Colour Doppler ultrasound unit – Aloka Prosound alpha-6
- High frequency linear array ultrasound transducer with a range 8, 12, 14 MHz.
- Ultrasound jelly.

Linear array transducer were preferred to sector transducer because of wider near field of view and the capability to combine high frequency gray scale and Colour Doppler images.

Doppler settings were standardized to compare the vascularity of thyroid pathologies among different patients and to ensure intra individual consistency. Observations were recorded in the proforma and hard copy of imaging mode.

Technique of scanning

Patients were scanned in supine position. Visualization was enhanced by performing the examination with the neck hyper extended and by asking the patient to swallow so as to elevate the subclavicular portion of the gland. A small pad was placed under the shoulders to provide better exposure of the neck, particularly in patients with short stocky habitus. Scans were obtained in standard transverse and longitudinal as well as multiple oblique positions. Alteration in echogenicity and echo texture were noted and focal nodules were localized, measured and characterized. Examination was laterally extended to include the region of carotid artery and jugular vein to identify enlarged jugular chain of lymph nodes, superiorly to visualize submandibular adenopathy and inferiorly to define any pathological supraclavicular lymph nodes.

Observations

The findings in the study on 100 patients referred to the Department of Radio diagnosis are given below.

Statistical significance of sex ratio in thyroid disorders is depicted in Table 1

| Sex | No. of Cases | Percentage |
|--------|--------------|------------|
| Male | 17 | 17% |
| Female | 83 | 83% |
| Total | 100 | 100% |

Table 1: Sex ratio of thyroid disorders.

The table shows that there were 83 females and 17 males of the total 100 cases. The male to female ratio in our study is 1:4.88. Patients were grouped into different age group according to their age, which is tabulated in Table 2

The age group studied included patients ranging from 75 years with youngest being female of 13 years and oldest male of 65 years. The major age group affected is between 20-40 years with 56 percentage cases.

| Age | No. of Cases | Percentage |
|-------|--------------|------------|
| 10-20 | 5 | 5% |
| 20-30 | 23 | 23% |
| 30-40 | 33 | 33% |
| 40-50 | 18 | 18% |
| > 50 | 21 | 21% |
| Total | 100 | 100% |

Table 2: Age Wise Distribution of the Cases.

The pattern of involvement of thyroid gland in various thyroid disorders is depicted in Table 3.

| Pattern of Involvement | No. of Cases | Percentage |
|------------------------|--------------|------------|
| Diffuse | 20 | 20% |
| Nodular | 80 | 80% |
| Total | 100 | 100% |

Table 3: Pattern of Involvement of Thyroid Disorders.

Thyroid disorders can either be diffuse or nodular. Thyroid disorders are predominantly nodular with ratio of diffuse to nodular being 1:4

The different sonographic criteria used to characterize diffuse thyroid disorders are classified in Table 4

| Sonographic Criteria | No. of Cases N (20) | Percentage |
|----------------------|------------------------|------------|
| Heterogeneity | 15 | 75% |
| Micronodulations | 3 | 15% |

Table 4: Sonographic Criteria in Diffuse Thyroid Diseases.

Heterogeneous involvement of thyroid parenchyma was a feature of 15 (75%) patients. Fine 1-6 mm micro nodules were present in only 3 (27%) patients.

Thyroid nodules were categorized into three types on the basis of echotexture pattern, which is shown in Table 5.

| Echotexture Pattern | No. of Cases | Percentage |
|---------------------|--------------|------------|
| Cystic | 15 | 18.75% |
| Solid | 41 | 51.25% |
| Mixed | 24 | 30% |
| Total | 80 | 100 |

Table 5: Incidence of Different Echotexture Pattern in Thyroid Nodules.

Out of total 80 thyroid nodules 18.75% were cystic, 51.25% were solid and 30% were mixed in echotexture.

Various B-Mode Criteria used in characterization of thyroid nodules are echogenicity, margins, halo and calcification. They are tabulated in Table 6.

| B-Mode Findings | No. of Cases N-49 | Percentage |
|---------------------|----------------------|------------|
| Echogenicity | | |
| Hypoechoic/Echofree | 44 | 55% |
| Isoechoic | 21 | 26.25% |
| Hyperechoic | 15 | 18.75% |
| HALO | | |
| Thin & Complete | 64 | 80% |
| Thick & Incomplete | 16 | 20% |
| MARGINS | | |
| Regular | 67 | 83.75% |
| Irregular | 13 | 16.25% |
| Calcification | | |
| Microcalcification | 5 | 6.25% |
| Coarse Or Egg Shell | 14 | 17.5% |

Table 6: Incidence of Different B-Mode Findings in Thyroid Nodules.

Out of the total 80 nodules 21 were Isoechoic, 15 Hyperchoic and 44 Hypoechoic. 64 patients have thin and complete halo around the nodule whereas 16 patients has thick, incomplete or absent halo. Margins of the nodule were regular in 67 cases whereas it was Irregular in 13 cases.

All the patients' didnot show calcification. Only 5 patients have micro calcification whereas 14 patients have evidence of coarse of eggshell calcification.

Nodular vascularity was determined by power Doppler and classified into 5 types which are shown in Table 7.

| Colour Doppler Pattern | No. of Cases | Percentage |
|------------------------|--------------|------------|
| Type I | 18 | 22.5% |
| Type II | 26 | 32.50% |
| Type III | 34 | 42.50% |
| Type IV | 2 | 2.5% |
| Total | 80 | 100% |

Table 7: Colour Doppler Pattern in Thyroid Nodules.

The Table shows that most of the patients have type II and III patterns Type I pattern in which there is only peripheral flow is seen in 18 (22.5%) nodules.

Type II pattern having peripheral and central component of flow with peripheral component predominating is seen in 26 (32.5%) patients. Type III pattern with central component predominating over the peripheral is seen in 34 (42.5%) patients. Only 2 patients have type IV pattern, which has only central flow.

Colour Doppler indices, which were taken into account in our study, were resistive index and peak systolic velocity. They are depicted in Table 8.

| Colour Doppler Indices | No. of Cases N-49 | Percentage |
|------------------------|-------------------|------------|
| Resistive Index (RI) | | |
| < 0.70 | 71 | 88.75% |
| > 0.70 | 9 | 11.25% |

Table 8: Colour Doppler Indices in Thyroid Nodules.

Most of the thyroid nodules (88.75 %) have resistive indices less than 0.7 with only 11.25 % having RI over 0.7.

On the basis of different B-mode criteria diagnosis was assigned to the patient, which was then compared with FNAC (fine needle aspiration cytology). Sensitivity and specificity of each diagnosis was calculated which is shown in Table 9

| Thyroid Pathologies | B-Mode | FNAC | False Positive | False Negative | Sensitivity | Specificity |
|-------------------------|--------|------|----------------|----------------|-------------|-------------|
| Simple Goitre | 11 | 12 | - | 1 | 92.3% | 100% |
| Solitary Thyroid Nodule | 15 | 14 | 3 | 2 | 85.7% | 95.6% |
| Multinodular Goitre | 43 | 40 | 3 | - | 100% | 93% |
| Adenoma | 15 | 14 | 4 | 5 | 73.6% | 94.2% |
| Carcinoma | 8 | 9 | 1 | 1 | 90% | 98.6% |
| Hashimoto's Thyroiditis | 4 | 5 | - | 1 | 80% | 100% |
| Graves disease | 4 | 6 | - | 2 | 75% | 100% |
| Total | 80 | 80 | 11 | 12 | 86.95% | - |

Table 9: Comparison of B-Mode Diagnosis with Fnac.

Sensitivity of B-mode to diagnose malignancy is good (90 %). Among diffuse thyroid diseases B-mode was able to diagnose correctly all but one case of simple goiter and sensitivity for Hashimoto's thyroiditis was 80 %. B-mode under diagnosed 2 cases of Grave's disease. Overall sensitivity of B-mode in diagnosing thyroid pathologies is 86.95 %.

The incidence of different B-mode and colour Doppler findings in benign and malignant thyroid nodules were studied retrospectively, after the diagnosis was confirmed on FNAC. These different findings are depicted in Table 10

| Ultrasound Findings | Benign | | Malignant | |
|---------------------|--------|--------|-----------|-------|
| | N | % | N | % |
| Echogenicity | | | | |
| Hypoechoic | 38 | 47.5% | 6 | 7.5% |
| Isoechoic | 19 | 23.75% | 2 | 2.5% |
| Hyperchoic | 14 | 17.5% | 1 | 1.25% |
| Halo | | | | |
| Thin & Complete | 62 | 77.5% | 1 | 1.25% |

| | | | | |
|------------------------|----|--------|---|-------|
| Thick & Incomplete | 9 | 11.25% | 8 | 10% |
| Margins | | | | |
| Regular | 64 | 80% | 2 | 2.5% |
| Irregular | 7 | 8.75% | 7 | 8.75% |
| Calcification | | | | |
| Microcalcification | 1 | 1.25% | 4 | 5% |
| Coarse or Egg Shell | 13 | 16.25% | 1 | 1.25% |
| Colour Doppler Pattern | | | | |
| Type I & II | 42 | 52.5% | 2 | 2.5% |
| Type III & IV | 29 | 36.25% | 7 | 8.75% |
| Resistive Index (RI) | | | | |
| < 0.75 | 68 | 85% | 1 | 1.25% |
| > 0.75 | 3 | 3.75% | 8 | 10% |

Table 10: B-Mode & Colour Doppler Features in Benign & Malignant Thyroid Nodules.

In our study most of the malignant lesions are hypo echoic with thick incomplete halo, irregular margins, micro calcification, resistive index > 0.70. Colour Doppler pattern type III & IV is 77.8 % sensitive and 59 % specific for malignancy.

Most of the benign lesions are isochoric or hyper echoic with thin complete peripheral halo, regular margins, coarse or eggshell calcification, resistive index < 0.70. Colour Doppler pattern type I & II is 59 % sensitive and 18.4 % specific for benign nodules.

The positive predictive value, negative predictive value and diagnostic accuracy of various gray scale and Doppler findings in differentiating malignant from benign lesions are shown in Table 11. The sensitivity and specificity of various gray scale and Doppler findings in differentiating malignant from benign lesions is depicted in the table below

| | PPV | NPV | Diagnostic Accuracy |
|--------------------------------|-------|-------|---------------------|
| Absent Halo | 47 % | 98 % | 87.5% |
| Irregular Margin | 50 % | 96.9% | 88.75% |
| Microcalcification | 80% | 92.8% | 21.25% |
| RI > 0.7 | 72.7% | 98.5% | 95% |
| Colour Doppler Type Iii and Iv | 19.4% | 95.4% | 61.25% |

Table 11

Thus micro calcification and RI Of 0.7 has high specificity in prediction of malignant thyroid nodules

The specificity of type III and IV flow in detection of malignancy is low.

The effect on sensitivity when type of flow is combined with gray scale features is shown in Table 13

Thus we see that on combination of colour Doppler and gray scale parameters the specificity is increased.

| | Carcinoma | Benign Nodules | Sensitivity | Specificity |
|--------------------|-----------|----------------|-------------|-------------|
| Absent halo | 8/9 | 9/71 | 88.8% | 87.3% |
| Irregular Margin | 7/9 | 7/71 | 77.7% | 90% |
| Microcalcification | 4/9 | 1/71 | 44.5% | 98.5% |
| RI > 0.7 | 8/9 | 3/71 | 88.8% | 95.7% |
| Type III & IV Flow | 7/9 | 29/71 | 77.7% | 59% |

Table 12

| | Carcinoma | Benign Nodules | Sensitivity | Specificity |
|--------------------------------------|-----------|----------------|-------------|-------------|
| Absent Halo +Type III & IV Alow | 6/9 | 6/71 | 66.7% | 91.5% |
| Irregular Margin +Type III & IV Flow | 5/9 | 4/71 | 55.5% | 94.36% |

Table 13

Discussion

Thyroid diseases diagnosis requires a holistic approach. Although the thyroid is a superficial organ easily accessible to physical examination, it needs various biochemical and radiological investigations to confirm the diagnosis.

The introduction of high-resolution ultrasonography had made it possible to detect many non-palpable nodules in the thyroid. The incidence of such lesions in the general population appears to be high approximately 10-40%.

The present study was concluded in a tertiary hospital from Oct. 2013-Sept 2014. The study population consisted of 100 patients who were diagnosed to have thyroid abnormality on ultrasonography.

In the present series, out of 100 patients, 83 were females and 17 were males. There is predominant involvement of thyroid gland in females with F: M ratio of 4.88:1.

Solbiati, *et al.* In their study of 401 patients found that 71.3% were females and 28.7% were males.

Khurana KK, *et al.* Reported their experience with 119 patients, out of which 100 (84%) were females and 19 (16%) males.

Summary and Conclusion

High resolution ultrasonography is highly sensitive in diagnosing thyroid disorders. Colour Doppler acts as an important adjunct to B-mode in increasing the accuracy.

In our study we concluded that -

- Females are more commonly affected than males.
- Most common age group affected is between 20-40 years.
- Nodular involvement of thyroid is more common than diffuse.
- Small hypo echoic 1-6 mm micro nodules are highly predictive of Hashimoto's thyroiditis
- Non-neoplastic nodules are more commonly hyper echoic or isoechoic whereas carcinomas are usually hypo echoic. Adenoma can present with variable echogenicity.
- Non-neoplastic nodules and adenomas usually present with regular margins whereas irregular margin is more a feature of carcinoma.

- Adenomas and non-neoplastic nodules are more commonly associated with thin and complete halo. Halo is thick, incomplete or absent in most of the carcinoma.
- Micro calcification has highest specificity for malignancy on B-mode.
- Coarse and eggshell calcification is highly predictive of benignity.
- Power Doppler pattern is not sensitivity and specific in differentiation of malignant and benign thyroid nodules in isolation.
- Resistive index < 0.70 in majority of non-neoplastic nodules. Carcinoma usually have RI > 0.70
- Specificity and positive predictive value of diagnosing a lesion as malignant increases, when combination of criteria is used, instead of individual criteria.

The goal of imaging should be to avoid extensive and costly evaluation in the most patients with benign disease without missing the minority of patients with thyroid cancer. High-resolution ultrasonography has proved high sensitivity in the detection of very small nodular lesions of the thyroid. Its specificity in definition of benign from malignant nature is being greatly improved by the new vascular pattern and resistive index parameters using pulsed and power Doppler. In this study we found no single criteria that could distinguish benign from malignant thyroid nodules with 100% reliability. FNAC (fine needle aspiration cytology) still offers the highest accuracy. Thyroid sonography is most useful in the differentiation between definitely benign nodules from suspicious which can then be subjected to further evaluation by FNAC (fine needle aspiration cytology).

References

1. Andre Van Harle., *et al.* "The thyroid nodule". *Annals of Internal Medicine* 96 (1982): 221-232.
2. Argalia G., *et al.* "Echo Doppler in the characterization of thyroid nodular disease". *Radiol Med (Torino)* 89.5 (1995): 651-657.
3. Ahuja A., *et al.* "Clinical significance of the comet-tail artifact in thyroid ultrasound". *Journal of Clinical Ultrasound* 24.3 (1996): 129-133.
4. Amodio F., *et al.* "Role of flowmetric analysis and of colour Doppler ultrasonography with contrast media in the different phases and follow up of Graves' disease". *Radiol Med (Torino)* 102.4 (2001): 233-237.
5. Andrej Lyschik., *et al.* "Diagnosis of thyroid cancer in Children: value of Gray-Scale and power Doppler US". *Radiology* 235.2 (2005): 604-613.
6. Black B M., *et al.* "Multicentricity of papillary adenocarcinoma of the thyroid: influence on treatment". *The Journal of Clinical Endocrinology & Metabolism* 20.1 (1960): 130-135.
7. Berghout A., *et al.* "Determination of thyroid volume as measured by ultrasonography in healthy adults in a non-iodine deficient area". *Clinical Endocrinology* 53.5 (1987): 273-280.

Submit your next manuscript to Scientia Ricerca Open Access and benefit from:

- Prompt and fair double blinded peer review from experts
- Fast and efficient online submission
- Timely updates about your manuscript status
- Sharing Option: Social Networking Enabled
- Open access: articles available free online
- Global attainment for your research

Submit your manuscript at:

<https://scientiaricerca.com/submit-manuscript.php>