

Serum Level of Anti-ccp in Patients with Active Pulmonary Tuberculosis

Sahar Emami*

Department of Veterinary, Shushtar Branch, Islamic Azad University, Shushtar, Khuzestan, Iran

*Corresponding Author: Sahar Emami, Department of Veterinary, Shushtar Branch, Islamic Azad University, Shushtar, Khuzestan, Iran.

Received: August 26, 2017; Published: September 06, 2017

Abstract

Background: Tuberculosis (TB) is a major public-health problem which principally affects the lungs, but may presents a variety of rheumatic manifestations, simulating rheumatologic disorders such as rheumatoid arthritis.

Methods: The main aim of this research was examined the anti-ccp level in 40 patient with active tuberculosis and 40 normal person's serum by ELISA.

Results: Mean levels of anti-CCP were significantly increased ($p < 0.01$) in patients with TB compared with controls: 29.8 (17.7) vs 10.8(7.9) U/ml, respectively.

Conclusion: Finally, a high percent of patient with active TB show an increased level of anti-ccp that could make difficulties to detecting TB Arthritis.

Key words: Tuberculosis; Cyclic citrullinated peptide antibody

Volume 1 Issue 2 September 2017

© All Copy rights are reserved by Sahar Emami.

Introduction

Tuberculosis (TB) is a major cause of human mortality and a worldwide public health problem. TB is a chronic, communicable disease caused by *M. tuberculosis* made distinctive by a destructive necrotizing granulomatous tissue response to the organisms. *Mycobacterium tuberculosis* causes two million deaths every year and every year 8 million of new cases of TB are diagnosed worldwide [1,2]. Both pulmonary TB and also cases of extra pulmonary disease are increasing [3]. About 10% of extra pulmonary TB affects the bones and joints [4]. Tuberculosis of bones and joints often presents as gradually worsening arthritis. Systemic and pulmonary symptoms are frequently absent, and the differential diagnosis must include other possible causes of septic osteoarticular disease, inflammatory arthritis. Osteoarticular tuberculosis rarely involves more than one joint, and this may help to differentiate it from other poly inflammatory diseases. Also, the radiographic changes of tuberculosis are slow to develop compared with those of pyogenic infections, and a reduction in the joint space is often a late occurrence [5]. Clinical confusion occurs especially when tuberculosis superinfects joints previously involved by inflammatory arthritis [6].

Citation: Sahar Emami. "Serum Level of Anti-ccp in Patients with Active Pulmonary Tuberculosis". *Pulmonary Research and Respiratory Care* 1.2 (2017): 26-30.

TB arthritis most commonly manifests as a mono arthritis of the weight bearing joints [7] but oligo articular or poly articular presentation is not rare and these findings may mimics inflammatory diseases such as the spondyloarthropathies [8] or rheumatoid arthritis (RA), or both [9]. Also, the serum of patients with TB may contain rheumatoid factors (RFs) in up to 40% of cases [10]. Patients with TB, even those without evidence of a direct musculoskeletal/local involvement, may present with a variety of rheumatic symptoms and signs including arthralgia and arthritis. The presence of rheumatic manifestations can lead to challenges in making or excluding a diagnosis of rheumatoid arthritis (RA) in patients infected with TB. For distinguishing between true RA and TB rheumatic symptoms, the presence of RF is of little help and can be confusing or misleading. Anti-cyclic citrullinated peptide (Anti-CCP) were detected in the serum samples of RA patients which shows high diagnostic accuracy. Citrulline is an unusual amino acid created by de-imitation of arginine residues in several proteins by the action of peptidyl arginine deaminase (PAD). The antibodies are detected by ELISA, where a synthetic cyclic citrullinated peptide (CCP) is used as substrate [11].

Materials and Methods

50 serum samples of patients with active lung tuberculosis (29 men and 11 woman) diagnosed according to clinical symptoms, radiologic studies, and laboratory confirmation including direct smear of sputum and sputum cultures for Mycobacterium tuberculosis and consecutively recruited from the department of tuberculosis and pulmonary disease in 5th Aboozar Hospital, in Ahvaz (Iran), an endemic area of tuberculosis Also 50 sex- and age-matched healthy subjects are studies for serum anti-ccp levels. Among the TB patients, the variables recorded were age, gender, time from diagnosis, duration of symptoms, fever, hemoptysis, cough, myalgia, and arthralgia and serum titer of anti-ccp. Anti-CCP antibodies were tested by commercial ELISA (Aeskulisa; Aesku. Diagnostics, Germany). Eachof the tests was performed and evaluated by operators whowere blinded to thepatients' clinical data, and radiological and microbiological results. The assay was performed according to the manufacturer's instructions. To determine the precision of the assay, the variability (intra and inter-assay) was assessed by examining its reproducibility on three serum samples selected to represent a range over the standard curve. Analytical sensitivity was 1.0 U/ml. Intra-and interassay CVs were 0.4-8.9% and 1.0-8.2%, respectively. All moral standards in this study were compiled and the project was approved by the University's ethics committee.

Statistical Analysis

Analysis was performed using the SPSS version 13, Student's t test and an χ^2 test to compare antibody titersor positivity rate, respectively, between patients with TB and controls. Pearson correlation coefficients were used to study the relationship between clinical measures and the level of anti-CCP. A value of $p < 0.01$ was considered significant.

Results

The patients (mean and SD 51.1+/-15.2 years; range, 19-81years) and healthy subjects (mean and SD 47.4+/-15.7 years; range, 23-85 years) ages were determined. The mean (SD) duration of TB related symptoms was 4.4 (1.7) months, 72% had fever, 78% had cough and 78%had hemoptysis. Rheumatic symptoms were relatively rare: arthralgia (2%), myalgia (4%), and eye and mouth dryness (1% and 3%, respectively). Mean (SD) levels of anti-CCP were significantly increased in patients with TB compared with controls: 29.8 (17.7) U/ml v 10.8 (7.9) U/ml ($p < 0.001$). Serum levels >15 U/ml were found in 17 (34%) patients compared with 2 (6%) controls ($p = 0.002$). Risk of pulmonary TB in subjects with anti-CCP serum level higher than 15 U/ml is almost 12.5 fold persons lower than the cut-off point (odds ratio = 12.4, CI 95%: 2.6-114.9). No relationship was found between the titer of anti-CCP and fever and any rheumatic symptom. Table 1 summarizes the demographic, clinical characteristics, and anti-CCP levels of patients with TB and healthy controls.

Characteristics	Patients with TB	Control Group
Age (years), mean (SD)	51.1 (15.2)	47.4 (15.7)
Sex (M/F)	29/21	30/20
Duration of symptoms (months), mean (SD)	4.4 (1.6)	-
Fever, No (%)	36 (72)	-
Cough, No (%)	39 (78)	-

Hemoptysis, No (%)	39 (78)	-
Rheumatologic symptoms, No (%)	6 (12)	-
Serum Anti-CCP level (> 15 IU)	17 (34)	3 (6)

Table 1: Demographic, clinical characteristics and anti-CCP levels of patients with TB and healthy controls.

Discussion

We sought to determine if anti-CCP was also associated with active pulmonary tuberculosis. The data presented here clearly indicate that anti-CCP can be found in the sera of unselected patients with TB. We found that 17 (34%) patients with TB have positive levels of anti-CCP. Although the presence of anti-CCP correlated with fever, it was not associated with symptoms and signs of arthritis.

Elkaya, *et al.* showed significantly increased anti-ccp levels in tuberculosis patients in comparison with controls. In this study, 32 percent of patients in comparison with 2.6 percent of controls show serum levels above the upper normal limits. They found correlation between increased serum anti-ccp and a history of prolonged of fever, as we noted in the present study. As we revealed, no correlation was found between anti-ccp level and rheumatic symptoms in Elkayam study [12].

Patients with TB may have a variety of rheumatic symptoms and signs, including arthralgia and arthritis and TB may imitate inflammatory rheumatic diseases, such as RA [13]. Also the radiological abnormalities observed in tuberculosis arthritis may simulate RA [14]. Antibodies to CCP are present in the sera of a majority of patients with RA, only rarely in the sera of patients with other diagnoses.

TB of the bones and joints, which constitutes a smaller part of extra pulmonary tuberculosis, is relatively rare in documented cases and can yield variable clinical manifestations. Many authors describe numerous cases of tuberculosis arthritis in patients presenting with monoarthritis [15-20]. Tuberculosis must always be considered in the differential diagnosis of chronic monoarthritis if devastating sequelae are to be avoided.

In recent years, various antibodies have been described in rheumatoid arthritis, and their clinical significance have been discussed. In particular, autoantibodies to citrullinated proteins such as fill grin and its circular form (cyclic citrullinate peptide: anti-ccp) are the most remarkable because of the reasonable sensitivity and high specificity in RA patients [21]. They are presenting most patients with RA and have been found to have a specificity of > 90% [22]. Although, anti-CCP may present in the serum of patients with psoriatic arthritis and have questioned its specificity [23].

The presence of anti-CCP in a few infectious diseases, as hepatitis C has been studied. Although, rheumatic factor (RF) is present in the great majority of patients with hepatitis C arthritis and in a proportion of patients with other sub-acute infections, particularly bacterial endocarditis, anti-CCP is negative in patients with HCV and measurement of anti-CCP may help in diagnosing RA in patients with chronic HCV infection [24].

Patients with active tuberculosis may have increased titers of various autoantibodies and serologic markers, such as rheumatic factor (RF), antinuclear antibody (ANA), anti cardiolipins, anti neutrophilic cytoplasmic antibody, anti-ds DNA, anti-Sm, anti-RNP and anti-Ro [25-27]. Genetic factors such as HLA and agene polymorphism of the citrullinating enzyme, peptidyl arginine deiminase, (that might express more stable mRNA and cause over-citrullination of proteins) might be associated with the breakage of self-tolerance and induction of autoimmunity against citrullinated proteins [20,28]. It is not clear whether the false positive anti-CCP reactivity seen in patients with TB is directed against citrullinated or non-citrullinated epitopes in the substrate for the CCP test [12].

Conclusion

A relatively high percentage of patients with active pulmonary tuberculosis show increased titers of anti-ccp which could impede diagnosis of articular TB. Therefore, it is recommended not to limit the diagnosis of arthritis types to anti-ccp test, and perform other assessments in order to differentiate articular TB.

Citation: Sahar Emami. "Serum Level of Anti-ccp in Patients with Active Pulmonary Tuberculosis". *Pulmonary Research and Respiratory Care* 1.2 (2017): 26-30.

References

1. World Health Organization. Global tuberculosis control: surveillance, planning, financing. WHO report 2005. Geneva, *World Health Organization* (WHO/HTM/TB/2005.349).
2. Corbett E., et al. "The growing burden of tuberculosis: global trends and interactions with the HIV epidemic". *Archives of Internal Medicine* 163 (2003): 1009-1021.
3. Dye C., et al. "Consensus statement. Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country, WHO Global Surveillance and Monitoring Project". *JAMA* 282.7 (1999): 677-86.
4. Malaviya AN and Kotwal PP. "Arthritis associated with tuberculosis". *Best Practice & Research: Clinical Rheumatology* 17 (2003): 319-343.
5. Hsu SH., et al. "Reappraisal of skeletal tuberculosis: role of radiological imaging". *Journal of the Formosan Medical Association* 92 (1993): 34-41.
6. Stecher DR., et al. "Tuberculous arthritis in the course of connective tissue disease: Report of 4 cases". *The Journal of Rheumatology* 19 (1992): 1418-20.
7. Watts HG and Lifeso RM. "Tuberculosis of bones and joints". *The Journal of Bone and Joint Surgery, American* 78 (1996): 288-298.
8. Hammoudeh M and Khanjar I. "Skeletal tuberculosis mimicking seronegative spondyloarthropathy". *Rheumatology International* 24 (2004): 50-52.
9. Bush DC and Schneider LH. "Tuberculosis of the hand and wrist". *Journal of Hand Surgery* 9 (1984): 391-398.
10. Isenberg DA., et al. "Profile of autoantibodies in the serum of patients with tuberculosis, klebsiella and other gram-negative infections". *Clinical & Experimental Immunology* 76 (1987): 516-523.
11. Schellekens GA., et al. "The diagnostic properties of rheumatoid arthritis antibodies recognizing a cyclic citrullinated peptide". *Arthritis & Rheumatology* 43 (2000): 155-163.
12. Elkayam O., et al. "Positive anti-cyclic citrullinated proteins and rheumatoid factor during active lung tuberculosis". *Annals of the Rheumatic Diseases* 65 (2006): 1110-1112.
13. Bush DC and Schneider LH. "Tuberculosis of the hand and wrist". *Journal of Hand Surgery* 9(1984): 391-398.
14. Hugosson C., et al. "Imaging of tuberculosis. Peripheral osteoarticular and soft tissue tuberculosis". *Acta Radiologica* 37 (1996): 512-516.
15. Al-Matar MJ., et al. "Isolated tuberculous monoarthritis mimicking oligoarticular juvenile rheumatoid arthritis". *The Journal of Rheumatology* 28.1 (2001): 204-206.
16. Hunfeld KP., et al. "Two cases of chronic arthritis of the forearm due to *Mycobacterium tuberculosis*". *European Journal of Clinical Microbiology & Infectious Diseases* 17.5(1998): 344-348.
17. Sawhney S and Murray KJ. "Isolated tuberculosis monoarthritis mimicking juvenile rheumatoid arthritis". *The Journal of Rheumatology* 29.4 (2002): 857-859.
18. Parra Parra I., "Osteoarticular tuberculosis of the right shoulder". *Archivos de Bronconeumología* 37.3 (2001): 154.
19. Martini M., et al. "Tuberculous osteitis". *Tunisie Medicale journal* 51.3 (1973):179-81.
20. Pernice W., et al. "Tuberculous arthritis--a rare, but important differential diagnosis in juvenile chronic arthritis". *Monatsschr Kinderheilkd* 139.4 (1991): 244-247.
21. Mimori T. "Clinical significance of anti-CCP antibodies in rheumatoid arthritis". *Internal Medicine* 44.11 (2005): 1122-1126.
22. Van Venrooij WJ., et al. "Anti-citrullinated protein/peptide antibody and its role in the diagnosis and prognosis of early rheumatoid arthritis". *Netherlands Journal of Medicine journal* 60 (2002): 383-388.
23. Vander Cruyssen B., et al. "Anti-citrullinated peptide antibodies may occur in patients with psoriatic arthritis". *Annals of the Rheumatic Diseases* 64 (2005): 1145-1149.
24. Wener MH., et al. "Absence of antibodies to cyclic citrullinated peptide in sera of patients with hepatitis C infection and cryoglobulinemia". *Arthritis & Rheumatology* 50 (2004): 2305-2308.
25. Adebajo AO., et al. "HazlemanBL.Autoantibodies in malaria, tuberculosis and hepatitis B in a west african population". *Clinical & Experimental Immunology* 93 (1993): 73-76.

26. Flores-Saurez LF, *et al.* "Prevalence of antinutrophilic cytoplasmic autoantibodies in patients with tuberculosis". *Rheumatology* 42 (2003): 711.
27. Elkayam O, *et al.* "Auto-antibody profiles in patients with active pulmonary tuberculosis". *International Journal of Tuberculosis and Lung Disease* 11.3 (2007): 306-310.
28. Suzuki A, *et al.* "Functional haplotypes of PADI4 encoding citrullinating enzyme peptidylarginine deaminase 4, are associated with reumatoid arthritis". *Nature Genetics* 34 (2003): 395-402.

Submit your next manuscript to Scientia Ricerca Open Access and benefit from:

- Prompt and fair double blinded peer review from experts
- Fast and efficient online submission
- Timely updates about your manuscript status
- Sharing Option: Social Networking Enabled
- Open access: articles available free online
- Global attainment for your research

Submit your manuscript at:

<https://scientiaricerca.com/submit-manuscript.php>